1. Introduction

The Greenland ice sheet is a vast body of ice covering approximately 80% of the surface of Greenland, and it is the second largest ice body in the world after the Antarctic ice sheet. Global warming has been affecting it severely because of the so-called polar amplification, i.e. the fact that changes in temperature resulting from greenhouse effects are larger than the average in geographical areas near the poles as in the case of Greenland. Specifically, the Greenland ice sheet has been melting and it is feared that if this process continues it will gradually reach a tipping point making it irreversible. For instance, Mouginot et al. (2019) calculated that there had been a sixfold increase in mass loss since the 1980s. Enderlin et al. (2014) provided evidence that surface melting rather than ice discharge was leading to sea level rises. Various other studies such as those by van der Broeke et al. (2017) and by the IMBIE Team (2020) similarly concluded that

methods

Nowadays, it is widely accepted that long memory is a property of many time series in the fields of hydrology, climatology, environmental studies, and economics and finance.

If one defines the spectral density function of a covariance stationary process as the Fourier transform of its autocovariances, i.e.,

$$f(\omega) = \sum_{j=-\infty}^{\infty} \gamma(u) e^{i\omega u} = \sum_{j=-\infty}^{\infty} \gamma(u) \cos(\omega u). \qquad (1)$$

where γ(u) ˜ Ը ϶ Գ ϵ ϶ ϵ

$$\int_{j_f}^{f} \left| J(u) \right| \qquad f \ .$$

coefficients in bold are those from the models selected on the basis of the statistical significance of the regressors, namely one with an intercept as well as a time trend in the case of NE, and one with an intercept only in the remaining cases.

TABLES 1 AND 2 ABOUT HERE

Table 2 reports the estimates of d as well as of the other coefficients. It can be seen that the estimated values of d are relatively high in all cases, ranging from 0.72 (NO) and 0.79 (SE) to 1.13 (SW). Evidence of mean

the UN Intergovernmental Panel on Climate Change (IPCC) concludes that “global warming will continue to increase in the near term (2021-2040) mainly due to increased cumulative CO_2 emissions in nearly all considered scenarios and modelled pathways. In the near term, global warming is more likely than not to reach 1.5°C even under the very low GHG emission scenario (SSP1-1.9) and likely or very likely to exceed 1.5°C under higher emissions scenarios. In other words, there is no evidence that current policies are succeeding in reducing carbon emissions, and therefore the expectation is that global warming will keep worsening. As a result, the most likely scenario is that the ice mass loss in Greenland will be reduced even further over time, as implied by our empirical results.

References

Bai, J. and P. Perron (2003), Bai, J., & Perron, P. (2003), Computation and analysis of multiple structural change models. Journal of Applied Econometrics, 18, 1-22.

Choi, Y., Morlighem, M., Rignot, E. and M. Wood (2021), Ice dynamics will remain a primary driver of Greenland ice sheet mass loss over the next century, Communications Earth & Environment, 2, article number 26.

Enderlin, E.M., Howat, I.M., Jeong, S., Noh, M.-J., van Angelen, J.H. and M. R. van den Broeke (2014), An improved mass budget for the Greenland ice sheet, Geophysical Research Letters, https://doi.org/10.1002/2013GL059010

Gil-Alana, L.A. (2008), Fractional integration and structural breaks at unknown periods of time, Journal of Time Series Analysis 29, 1, 163-185.

Granger, C.W.J. (1980), Long memory relationships and the aggregation of dynamic models, Journal of Econometrics 14, 2, 227-238.

Granger, C.W.J. and R. Joyeux (1980), An Introduction to Long Memory Time Series Models and Fractional Differencing, Journal of Time Series Analysis 1, 1, 15-29.

Hosking, J.R.M. (1981) Fractional differencing.Biometrika, 68, 168–176.

Hurst, H.E. (1956), The Problem of Long-Term Storage in Reservoirs, Hydrology Science Journal 1, 13–27.

IMBIE Team (2020), Mass balance of the Greenland Ice Sheet from 1992 to 2018, Nature, 579, 233–239.

IPCC (Intergovernmental Panel on Climate Change) (2023), "Climate Change 2023 – Synthesis Report", United Nations, https://www.ipcc.ch/report/ar6/syr/

Khan, S.A., Aschwanden, A., Bjørk, A.A., Wahr, J.,Kjeldsen, K.K. and K.H. Kjær (2015), Greenland ice sheet mass balance: a review, Reports on Progress in Physics, 78, 046801

Mandelbrot, B., (1971), A Fast Fractional Gaussian Noise Generator, Water Resources Research 7, 3, 543-553.

Mandelbrot, B. and J.R. Wallis (1971), Computer Experiments With Fractional Gaussian Noises: Part 1, Averages and Variances, Water Resources Research 5, 1, 228-241.

Mankoff, K.D., Solgaard, A., Colgan, W., Ahlstrøm, A.P., Khan, S.A., and Fausto, R.S. (2020). Greenland Ice Sheet solid ice discharge from 1986 through March 2020, Earth System Science Data, 12, 1367–1383.

Mouginot, J., Rignot, E., Bjørk, A.A. and M. Wood (2019), Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018, Earth, Atmospheric, and Planetary Sciences, https://doi.org/10.1073/pnas.1904242116

Table 1: Estimates of the integration order, d

Series	No deterministic terms	An intercept	An intercept and a linear time trend
CE	0.98 (0.92, 1.06)	**0.82 (0.74, 0.93)**	0.82 (0.74, 0.93)
CW	1.02 (0.96, 1.11)	**0.93 (0.85, 1.02)**	0.93 (0.85, 1.02)
NE	0.97 (0.90, 1.04)	0.82 (0.75, 0.90)	**0.82 (0.74, 0.90)**
NO	0.97 (0.91, 1.05)	**0.72 (0.64, 0.82)**	0.72 (0.64, 0.83)
NW	1.00 (0.94, 1.07)	**0.84 (0.77, 0.93)**	0.84 (0.78, 0.93)
SE	0.99 (0.93, 1.06)	**0.79 (0.74, 0.85)**	0.79 (0.74, 0.85)
SW	1.01 (0.95, 1.10)	**1.13 (1.04, 1.24)**	1.13 (1.04, 1.24)

In parenthesis the 95% confidence bands for the estimates of d. Values in bold are from the selected specification on the basis of the statistical significance of the regressors for each series.

Table 2: Estimated coefficients for the selected models

Series	No deterministic terms	An intercept	An intercept and a linear time trend	Seasonality

Table 4: Estimated coefficients before and after the breaks

Series	No deterministic terms	An intercept	An intercept and a linear time trend	Seasonality
i) First sub-samples				
CE	0.73 (0.56, 0.99)	67.8587 (28.33)	---	0.143
CW	0.57 (0.43, 0.81)	71.2801 (40.92)	---	0.078
NE	0.76 (0.67, 0.88)	22.7247 (37.64)	---	0.044
NO	0.59 (0.46, 0.76)	23.2706 (29.62)	---	0.011
NW	0.81 (0.71, 0.94)	93.8201 (55-66)	---	-0.067
SE	0.66 (0.57, 0.77)	143.1049 (58.05)	---	-0.027
SW	0.96 (0.83, 1.14)	20.5364 (45.84)	---	-0.004
i) Second sub-samples				
CE	0.93 (0.85, 1.03)	74.4772 (82.36)	---	-0.113
CW	1.11 (1.01, 1.22)	77.7730 (67.08)	---	0.017
NE	1.17 (1.06, 1.31)	26.5309 (37.64)	---	-0.083
NO	1.31 (1.18, 1.47)	24.9734 (29.62)	---	-0.0006
NW	0.87 (0.77, 1.00)	103.8391 (55.66)	---	-0.034
SE	0.91 (0.84, 1.00)	137.1984 (58.05)	---	-0.010
SW	1.22 (1.11, 1.35)	19.7333 (45.84)	---	-0.074