Department of Economics and Finance

Working Paper No. 24-01

Economics and Finance Working Paper Series

Guglielmo Maria Caporale and Luis Alberiko Gil-Alana

A Long-Memory Model for Multiple Cycles with an Application to the S&P500

February 2024

http://www.brunel.ac.uk/economics

A LONG-MEMORY MODEL FOR MULTIPLE CYCLES WITH AN APPLICATION TO THE S&P500

Guglielmo Maria Caporale, Brunel University London, UK

where -1),

d_j is another real value corresponding to the order of integration of the cycle that explodes (i.e., it goes to infinity) ; m stands for

the estimation of as many parameters as in the case of ARMA models. In addition, Bloomfield (1973) model is stationary across all its values (see Gil-Alana, 2004).

In the empirical application carried out below we assume that p = 4, and that w(r;1) = 0, so the first cyclical component corresponds to the long run or zero frequency. In such a case, the summand becomes , which can be expressed as , with the pole or singularity in the spectrum going to infinity at the zero frequency (Granger, 1980, Granger and Joyeux, 1980 and Hosking, 1981). For the other two cyclical structures, we choose the frequencies on the basis of the values of

this test against various alternatives. A similar experiment was conducted in Gil-Alana (2001), though in that case m = 1 and thus only a single cyclical structure was allowed. By contrast, in the present study we assume that m = 3 and that the Data Generating Process (DGP) is characterised by the following orders of integration: d1= 0.75, d2 = 0.50 and d3 = 0.25; this implies that the first cyclical structure is highly persistent and non-stationary, the second one is on the borderline between the stationary and nonstationary case, and the third one is stationary. For the length of the cycles we impose j = 10, 100 and 250, with different sample sizes. These values are arbitrary, though the results were found to be robust to choosing other values. For the alternative hypotheses we consider values for the three orders of integration of 0.25, 0.50 and 0.75. For j we choose the same values as in the true model, therefore the size of the test is reported in the tables for d = $(0.75, 0.50, 0.25)^T$.

INSERT TABLES 1, 2 AND 3 ABOUT HERE

Table 1

log transformation in the upper panel, and their corresponding periodograms in the lower one. Both exhibit a very large value at the smallest (zero) frequency, which

(11)

and make two

Ng, S. and Perron, P. (2001), Lag length selection and the Construction of Unit Root Tests with Good Size and Power. Econometrica, 69, 1519-1554.

Robinson, P.M. (1994) Efficient tests of nonstationary hypotheses, Journal of the American Statistical Association 89, 1420-1437.

Sadek, N. and A. Khotanzad (2004) K-factor Gegenbauer ARMA process for network traffic simulation, Computers and Communications 2, 963-968. Schmdit, P. and P.C.B.

Phillips (1992), LM Tests for a Unit Root in the Presence of Deterministic Trends, Oxford Bulletin of Economics and Statistics 54, 257-287.

Figure 1: Data in levels and periodograms

Figure 2: Data in first differences and periodograms

Original data	Logged transformed data
Periodogram original data	Periodogram logged
Periodogram original data	Periodogram logged
	

The values on the horizontal axis correspond

Table 6: Frequencies with the highest values at

(1 L) Data			(1 L) Log data		
j	T / j	Value at Periodogram	J	T / j	Value at Periodogram